

RESEARCH METHODS & STATISTICS

Evaluation of the Impact of Implementing the Emergency Medical Services Traumatic Brain Injury Guidelines in Arizona: The Excellence in Prehospital Injury Care (EPIC) Study Methodology

Daniel W. Spaite, MD, Bentley J. Bobrow, MD, Uwe Stolz, PhD, MPH, Duane Sherrill, PhD, Vatsal Chikani, MPH, Bruce Barnhart, RN, Michael Sotelo, Joshua B. Gaither, MD, Chad Viscusi, MD, P. David Adelson, MD, and Kurt R. Denninghoff, MD

Abstract

Traumatic brain injury (TBI) exacts a great toll on society. Fortunately, there is growing evidence that the management of TBI in the early minutes after injury may significantly reduce morbidity and mortality. In response, evidence-based prehospital and in-hospital TBI treatment guidelines have been established by authoritative bodies. However, no large studies have yet evaluated the effectiveness of implementing these guidelines in the prehospital setting. This article describes the background, design, implementation, emergency medical services (EMS) treatment protocols, and statistical analysis of a prospective, controlled (before/after), statewide study designed to evaluate the effect of implementing the EMS TBI guidelines—the Excellence in Prehospital Injury Care (EPIC) study (NIH/NINDS R01NS071049, “EPIC”; and 3R01NS071049-S1, “EPIC4Kids”).

The specific aim of the study is to test the hypothesis that statewide implementation of the international adult and pediatric EMS TBI guidelines will significantly reduce mortality and improve nonmortality outcomes in patients with moderate or severe TBI. Furthermore, it will specifically evaluate the effect of guideline implementation on outcomes in the subgroup of patients who are intubated in the field. Over the course of the entire study (~9 years), it is estimated that approximately 25,000 patients will be enrolled.

ACADEMIC EMERGENCY MEDICINE 2014;21:818-830 © 2014 by the Society for Academic Emergency Medicine

It is difficult to overstate the societal impact of traumatic brain injury (TBI). Annually in the United States, nearly 1.7 million victims of TBI are seen in emergency department (EDs), 275,000 are hospitalized, and over 50,000 die.¹⁻³ The Centers for Disease Control and Prevention (CDC) estimates that at least 5.3 million

From the Arizona Emergency Medicine Research Center (DWS, BJB, US, VC, BB, MS, JBG, CV, KRD), the Department of Emergency Medicine (DWS, US, BB, JBG, KRD), Barrow Neurological Institute at Phoenix Children’s Hospital and the Department of Child Health/Neurosurgery (PDA), College of Medicine, The University of Arizona, Tucson, AZ; the Arizona Department of Health Services (BJB, VC), Phoenix, AZ; the Maricopa Integrated Health System (BJB), Phoenix, AZ; and the College of Public Health, University of Arizona (DS), Tucson, AZ.

Received October 16, 2013; revision received February 18, 2014; accepted February 28, 2014.

Presented at the 9th World Congress on Brain Injury of the International Brain Injury Association, Edinburgh, Scotland, March 2012.

Research reported in this publication was supported by the National Institute of Neurological Disorders and Stroke of the National Institutes of Health under Award R01NS071049. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

This is a methods paper and not the clinical study. However, the study itself is registered at ClinicalTrials.gov, NCT01339702.

The authors have no potential conflicts of interest to disclose.

Supervising Editor: Craig D. Newgard, MD, MPH.

Address for correspondence and reprints: Daniel W. Spaite, MD; e-mail: dan@aemrc.arizona.edu.

Americans (about 2% of the population) have TBI-related long-term requirements for help performing activities of daily living.⁴ The direct and indirect costs of TBI totaled an estimated \$60 billion in 2000.² While the burden of injury is enormous across all age groups, children are disproportionately affected.⁵ In fact, TBI is the leading cause of death and disability in children.⁶

In major TBI (the term being used here for both moderate and severe injury), after the initial trauma (“primary brain injury”), additional secondary injury often occurs soon after the event.^{1,7–11} There is growing evidence that the care provided in the first few minutes may have a significant effect on outcome. Failure to immediately optimize treatment and limit secondary injury may lead to neurologic damage that is indelible, despite subsequent heroic in-hospital efforts.^{6–18} Based on these findings, evidence-based guidelines have been promulgated for in-hospital^{6,7,19} and prehospital^{10,19} TBI treatment.

The emergency medical services (EMS) TBI guidelines focus on the prevention and rapid correction of hypotension, hypoxia, and hyperventilation (in patients receiving positive-pressure ventilation). This is because of a large and growing literature showing the disastrous effect of these factors on outcomes.

Hypotension is common during both EMS and in-hospital TBI care^{8,20,21} and significantly decreases survival.^{6,8,9,20–38} A single episode is associated with a doubling of mortality, and this risk increases dramatically with repeated episodes (odds ratio = 8.1 in one study).²⁷ Conversely, the aggressive correction of hypotension early in the course of TBI is associated with improved outcomes.^{6,7,10,22,39}

Hypoxia (O₂ saturation < 90%) occurs frequently during the prehospital care of TBI patients^{13,16,31,40,41} and is associated with a major increase in mortality,^{8,20,22,24,28–31,42–44} with some studies reporting as much as a sevenfold increase.²⁸ Even a single hypoxic measurement in the field is associated with increased risk of death.^{8,42,43}

Until the early 1990s, “therapeutic hyperventilation” for TBI was commonly used because of studies suggesting it might improve outcome.^{45–49} However, there is now overwhelming evidence that even mild to moderate hyperventilation is detrimental to the brain due to vasoconstriction-induced ischemia and other factors.^{3,8,10,14,23,50–76} Unfortunately, inadvertent hyperventilation is extremely common in EMS when intubated patients are manually ventilated.^{3,14,51,52,77–84} The resulting hypocapnea is often severe, with end-tidal carbon dioxide (ETCO₂) levels of <25 mm Hg in as many as one-third of cases and <30 mm Hg in two-thirds of cases.⁷⁸ Denninghoff et al.⁸⁵ identified that, in the acute setting, failure to properly control postintubation ventilation is associated with a sixfold increase in severity-adjusted mortality.

The foregoing discussion highlights why the proper management of oxygenation, ventilation, and hemodynamics are at the core of the TBI guidelines. This robust literature, along with studies showing the effect of implementing the in-hospital guidelines,^{2,86–89} make it likely that implementing the guidelines in the field will significantly improve outcomes. However, the guideline development process acknowledged that the evidence supporting the EMS TBI treatment recommendations is

weak,^{6,7,10,19} and no large studies have yet evaluated their effectiveness. Here we describe a large prospective, before–after, statewide study evaluating the effect of implementing the EMS TBI guidelines—the Excellence in Prehospital Injury Care (EPIC) study. EPIC is funded by the National Institutes of Health (NIH/NINDS R01NS071049, “EPIC”; and 3R01NS071049-S1, “EPIC4K-ids”; ClinicalTrials.gov NCT01339702).

It is important to note from the outset that EPIC is an effectiveness study. It is not designed to identify the efficacy of specific, individual interventions. Rather, it is evaluating the effectiveness of the entire guideline “bundle” in a “real-world” setting across a vast demography and geography. This type of outcomes research goes hand-in-hand with classical randomized clinical trials aimed at identifying the efficacy of highly defined, carefully constructed, specific interventional questions.^{90,91}

The objectives of the study are to evaluate the following hypotheses:

- *Overall Study Hypothesis*—statewide implementation of the adult and pediatric EMS TBI guidelines will significantly reduce mortality and improve nonmortality outcomes in patients with moderate and severe TBI.
- *Specific Hypotheses*—implementation of the guidelines will: 1) significantly reduce overall mortality, 2) significantly reduce mortality among patients who are intubated in the field, and 3) significantly improve nonmortality outcomes (e.g., hospital length-of-stay, patient disposition).

Preimplementation TBI Care in Arizona

A prestudy survey evaluating 51 agencies (covering about 75% of the population) demonstrated highly variable TBI treatment across the state of Arizona. Only half had protocols specifying target ranges for O₂ saturation or blood pressure (BP), and only one-third had specific treatment protocols. While nearly 70% of agencies had ETCO₂ monitoring capabilities, only one-third of these had protocols specifying postintubation ETCO₂ targets. No agency had implemented the guidelines, thus creating the opportunity to prospectively evaluate the effect of implementation.

METHODS

Study Design

The EPIC study is a prospective, statewide, before–after, system study^{91–99} evaluating implementation of the EMS TBI guidelines throughout Arizona. EPIC has three phases (Figure 1). Phase 1 was the preimplementation (“before”) phase during which baseline risk adjustment and outcome measures were collected, from 2007 until implementation (most agencies implemented in 2012). Phase 2 was a “run-in” period. For each agency, Phase 2 began at the initiation of training and continued until implementation is complete. Data from Phase 2 are excluded from the analysis. Phase 3 is the postimplementation (“after”) phase. A study timeline is given in Figure 2.

Ethical Considerations

The necessary regulatory approvals for the EPIC project have been obtained from the Arizona Department of

Table 1
Specific, Guideline-based Management EMS Protocols

Age definitions* for monitoring and management ^{6,19}
<p>Infant: 0 to 24 months Child: 2 to 14 years Adolescent: 15 to 17 years Adult: 18 years and older</p>
<p>Management of airway/oxygenation Prevention, identification, and treatment of hypoxia (O₂ saturation < 90% and/or cyanosis) is a major emphasis.^{6-8,10,19,20,22,44} Care is initiated by continuous high-flow O₂ for all <i>potential</i> TBI cases. If high-flow O₂ fails to correct hypoxia, airway repositioning maneuvers are performed. If this does not restore eupoxia, or if there is inadequate ventilatory effort, bag-valve-mask (BVM) ventilation is performed using appropriate airway adjuncts (e.g., oropharyngeal airway). If airway compromise or hypoxia persists, ETI is performed if an experienced ALS provider is available.^{6-8,10,11,19,40,140} Following ETI, tube placement is confirmed via multiple means including ETCO₂ detection/capnography.</p>
<p>Management of ventilation Special emphasis is placed on identifying and treating hypoventilation as well as preventing hyperventilation when assisting ventilation. <i>Hypoventilation</i> (ineffective respiratory rate for age, shallow or irregular respirations, periods of apnea, or measured hypercarbia): if there is evidence of hypoventilation despite high-flow O₂ therapy, assisted ventilation is initiated via BVM and, if ineffective, ETI is performed.^{6,7,10,19,51,52,140} <i>Intubated patients</i>: ETCO₂ target is 40 mm Hg with careful attempts to maintain ETCO₂ between 35 and 45 mm Hg when monitoring is available.^{6,7,10,14,15,52} All agencies are encouraged to use ventilation rate timers and flow-controlled ventilation bags to assist in maintaining age-appropriate ventilatory rates and decrease the risk of inadvertent hyperventilation.^{6,7,10,51,52,77,78,80-84} Target initial ventilatory rates are from the pediatric and prehospital TBI guidelines:^{6,10,19} *infants, 25 breaths/min; children, 20 breaths/min; adolescents/adults, 10 breaths/min. <i>Mechanical ventilation</i>: When available, ventilators will be used postintubation to optimize ventilatory parameters and O₂ therapy.^{6,17,51,52,141,142} Target tidal volume will be 6 to 7 mL/kg with rates adjusted to keep the ETCO₂ within target range. This is consistent with the TBI guidelines and the recent literature showing that intrathoracic pressure, lung mechanics, hemodynamics, and intracranial pressure are optimized by this tidal volume compared to the "classic" 10 to 12 mL/kg that remains common in EMS and other settings.^{51,77,142-147} <i>Nonintubated patients</i>: All relevant monitoring/treatment will be applied except ETCO₂ monitoring. <i>Ventilation for impending cerebral herniation</i>: During <i>early</i> TBI management (prehospital/ED), the guidelines recommend strongly against any "prophylactic hyperventilation" even with severe brain injury, because there is strong evidence that this is detrimental. The guidelines recommend hyperventilation <i>only</i> for <i>obvious</i> signs of herniation. In such cases, only <i>mild</i> hyperventilation is recommended:^{6,7,10,19} *infants, 30 breaths/min; children, 25 breaths/min; adolescents and adults, 20 breaths/min. The ETCO₂ target range during "mild therapeutic hyperventilation" is 30 to 35 mm Hg.</p>
<p>Management of blood pressure <i>Hypotension</i>: Most TBI studies have used sBP < 90 mm Hg to identify hypotension in adults and older children.^{8,20-23,25-27,30-38} In the pediatric guidelines, hypotension is defined as sBP below the fifth percentile for age. This is estimated using the formula [70 mm Hg + (age × 2)]^{6,10,19,148} Age-appropriate definitions for hypotension Adults: < 90 mm Hg Adolescents: < 90 mm Hg Children age ≥10 years: < 90 mm Hg Infants and children ages younger than 10 years: 70 mm Hg + (age × 2) Adults and children age 10 years and older <i>Treatment of hypotension</i>: The TBI guidelines recommend keeping the sBP ≥ 90 mm Hg.^{6,7,10,19} A single sBP measurement < 90 mm Hg will initiate intravenous (IV) fluid resuscitation with an initial bolus of 1 L of normal saline or Ringer's lactate in adults and adolescents and 20 mL/kg in older children. This will be followed by IV fluid administration at sufficient rate and volume to keep sBP ≥ 90 mm Hg.^{6,7,10,19} <i>Treatment of hypertension</i>: In TBI, treatment of acute hypertension is not recommended.^{6,7,10,19,149} However, IV fluids will be restricted to a minimal "keep open" rate in patients with sBP ≥ 140 mm Hg. Infants and children age younger than 10 years <i>Treatment of hypotension</i>: For hypotension or other signs of shock, isotonic fluids will be given. Sufficient volume (via 20 mL/kg boluses every 5 minutes) will be given to return sBP to at least [70 mm Hg + (age × 2)].^{6,10,19} Intraosseous access should be attempted if there is hypotension or other signs of shock and peripheral venous access cannot be quickly established. <i>Treatment of hypertension</i>: As with adults and older children, treatment of acute hypertension is not recommended in TBI.^{6,7,10,19,149} IV fluids will be restricted to a minimal keep open rate in infants/young children with sBP ≥ 100 mm Hg.</p>
<p>ETCO₂ = end-tidal carbon dioxide; ETI = endotracheal intubation; TBI = traumatic brain injury.</p>

treatment protocols. Ongoing feedback to the agencies occurs through updates on progress of the study via multiple means and venues to providers, agency leaders, and medical directors.

Outcome and Severity/Risk Adjustment Measures. Using the Arizona State Trauma Registry

(ASTR) and the EMS patient care records, EPIC has access to an extensive array of risk adjustment and outcome measures (Table 2; Data Supplements S1 and S2, available as supporting information in the online version of this paper).^{91,94,108,109} The risk adjustment measures are used for inclusion or exclusion from the study and for risk stratification and analysis. The primary

Table 2
Severity/Risk Adjustment and Outcome Measures

<p>Risk adjustment/severity measures^{91,108,109} (from EMS patient care reports and the ASTR) <i>Demographics/history/EMS system response:</i> Age, sex, mechanism of injury (ICD-9/10-CM E-code Injury Descriptors), preexisting medical conditions, EMS time intervals¹⁵⁰ (e.g., interval from 9-1-1 call to trauma center arrival). <i>Prehospital data:</i> Initial and additional respiratory rate; heart rate; sBP; Glasgow Coma Scale (GCS) score; O₂ saturation and ETCO₂; prehospital procedures including IV fluids given (mL), ETI, rapid sequence intubation, supraglottic airway use, RTS, prehospital impression (conditions/"diagnoses") by EMS. <i>Trauma center data:</i> The ASTR data set contains nearly 200 data elements related to demography, severity, risk adjustment, treatment, procedures, and outcomes. Examples (detailed data set in Data Supplement S3)—initial and additional vital signs, GCS, RTS, probability of survival (TRISS), procedures (ICD-9/10-CM procedure codes), need for emergent surgery, need for emergent neurosurgery, intracranial pressure monitoring (yes/no), Abbreviated Injury Scale score, ISS, injury diagnoses (ICD-9/10-CM N-codes), preexisting comorbidities, hospital complications, hospital/ICU length of stay.</p> <p>Outcome measures^{7,8,10,11,17,18,23,85,86,95,96} <i>Primary outcome measure:</i> Survival to hospital discharge. <i>Secondary outcomes:</i> Prehospital mortality (patients with spontaneous circulation on EMS arrival who then require cardiopulmonary resuscitation by EMS and have no signs of life on arrival at the trauma center), hospital days, ICU days, total ventilator days, hospital complications, discharge disposition (e.g., home, long-term care facility, inpatient rehabilitation), and trauma center costs (charges).</p>
<p>ASTR = Arizona State Trauma Registry; ETCO₂ = end-tidal carbon dioxide; ETI = endotracheal intubation; ICU = intensive care unit; ISS = Injury Severity Score; RTS = Revised Trauma Score; TRISS = Trauma and Injury Severity Score.</p>

outcome measure is survival to hospital discharge.^{7,8,10,11,17,18,23,85,86,89,95,96} The secondary outcomes are prehospital mortality, hospital days, intensive care unit (ICU) days, ventilator days, hospital complications, discharge disposition (e.g., home, long-term care facility), and trauma center costs (charges).^{86-89,91}

Data Collection. Because EPIC is an ADHS-sponsored public health initiative, the State Attorney General has determined that it is exempt from HIPAA policy. The ADHS has maintained the ASTR since 2005 (with complete data since 2007). There is a detailed and explicit set of inclusion and exclusion criteria that are used by all trauma centers to determine which cases are submitted to the ASTR. Cases meeting specific EPIC study inclusion criteria are identified through the ASTR. This allows development of agency-specific lists of EPIC cases for linking the EMS patient care records to the ASTR data.

Prehospital/Trauma Center Case Linkage. Approximately 65% of EMS cases in Arizona are cared for by agencies that use paper-based systems, and the remainder use electronic patient care records. In agencies with paper-based records or with electronic patient care records where study staff do not have direct, online electronic access, the agency-specific case list is used to identify the matching patient care records by hand, and then study staff verify the linkage. Because of the high quality of the ASTR demographic data, in over 90% of cases the patient care record is easily matched by using the five "base" identifiers to establish patient identity (first name, last name, date of birth, sex, and incident date). If an unequivocal match is not made using the base identifiers, study staff review other fields based on incident street location, incident type or mechanism (e.g., motor vehicle crash), incident time, and specific EMS and trauma center case descriptions (e.g. specific descriptions of external injuries). Such detailed searching for linkage occurs in fewer than 10% of cases. We have direct, online access to electronic patient care records

for approximately 35% of study cases. The same matching process is followed for these as for paper-based agencies. Upon confirmation of a matched record, a unique identifier is established to merge the prehospital and hospital record for all future reference queries.

To demonstrate our ability to link the data, we performed a pilot study. Using the ASTR, we identified 75 consecutive severe TBI patients being brought to a trauma center by EMS and were able to match 100% of the cases and to verify that the EMS and trauma center data were available for all 75.

The EPIC electronic data warehouse is the repository for all cases meeting inclusion criteria. The trauma center data are stored on MS SQL 2008 servers, and the EMS data are stored on MY SQL 5.5 servers. Both data sets are stored on encrypted disk drives and transmitted only via encrypted file transfer or encrypted e-mail. A secure, encrypted key is used for linkage between the two data sets (Data Supplements S1 and S2).

Recent studies have identified that, in some systems, seriously injured trauma patients may be taken to non-trauma hospitals.¹¹⁰⁻¹¹² However, we believe that the statewide trauma system in Arizona creates a very high case capture rate for EPIC, since nearly all major TBI patients go to trauma centers (either directly or by transfer). To verify this, we used the comprehensive data from the ADHS Statewide Hospital Discharge and ED database. Because the primary outcome in the EPIC study is hospital mortality, we assessed the TBI deaths occurring in 2008. There were only 32 TBI deaths in the entire state that stayed at nontrauma centers (and, thus, were not in the ASTR). Essentially all of these were elderly patients with nonsurvivable injuries who were kept in local hospitals for supportive care. Because we have ongoing access to the discharge database, we will be able to identify TBI deaths occurring at nontrauma centers throughout the course of the study to help assess the effect of this factor on the case capture rate.

Evaluation of Guideline Compliance. The EPIC data elements deal specifically with guideline-related issues.

The data extraction process from the patient care records identifies guideline-relevant information (e.g., O₂ saturation, treating with high-flow O₂, sBP, IV fluids infused). These will be used to evaluate guideline compliance.

Sample Size

The study was powered based upon the EMS-intubated TBI subgroup because sufficient literature exists to make a reasonable estimate of treatment effect size in this cohort. First, using ASTR data (2007 through 2009), we are able to estimate the number of intubated cases that will be enrolled in Phase 1 (about 4¾ years in length, including 1,391 adults and 318 children [age < 18 years]).

Because no human studies have directly evaluated the effect of the EMS guidelines, the best data available for estimating the potential benefit (and, therefore, the number of cases needed in Phase 3) come from Wang et al.,¹⁰⁴ Davis et al.,¹⁷ and Denninghoff et al.⁸⁵ In these studies, intubated TBI patients transported with controlled ETCO₂ levels were compared to patients who had manual ventilation. These studies showed a remarkable 42% to 66% relative decrease in mortality in the properly ventilated cohorts. Based on these studies, a reasonable “best-case” estimate of treatment effect is a 42% relative reduction in mortality in intubated patients.

The study is powered based on an a priori plan of performing two sequential analyses. To calculate the power of the first interim analysis to detect a significant difference in mortality, the statistical “best-case” scenario was used (i.e., 42% relative decrease in mortality, an absolute decrease from 42.6% to 24.7% in adults, an absolute reduction from 36.9% to 21.4% in children, and a cluster effect of zero). Given these estimates for the interim analysis, in adults we will have a power of 0.99 and in children 0.77 ($\alpha = 0.01$, two-tailed) to detect a significant reduction in mortality (Fisher’s exact test).^{113–119}

The final analysis was powered based on a conservative treatment effect (21% relative decrease in mortality for adults and 34% decrease in children) and the assumption that there will be a significant clustering effect. The adult final analysis assumptions are as follows: Phase 1 intubated cases 1,391; Phase 3 intubated cases 987; $\alpha = 0.04$; intraclass correlation = 0.01; design effect = 2.184. Given these assumptions, the power of the study to detect a 21% relative reduction in mortality is 0.79.

Given the above calculations, we are able to estimate the number of moderate and severe, nonintubated TBI cases that will be enrolled. This is an important cohort to evaluate because previous inpatient studies make it likely that this group will benefit from guideline treatment as well.^{86–89} In adults, we estimate about 10,305 cases in Phase 1 and about 8,015 in Phase 3 (about 18,320 combined). Thus, in total, we expect to enroll over 20,000 adults in the EPIC study (2,416 intubated cases + 18,320 severe, nonintubated cases, totaling 20,736). In children, we anticipate enrolling 2,389 cases in Phase 1 and 1,760 in Phase 3 for a total of 4,149 cases of intubated and nonintubated TBI. Thus, the entire EPIC study is expected to enroll approximately 25,000

cases. Data Supplements S3 and S4 (available as supporting information in the online version of this paper) contain detailed descriptions of the analysis and sample size estimations.

Data Analysis

Descriptive statistics will be reported as means and standard deviations when data are normally distributed and median and interquartile ranges or 95% confidence intervals as appropriate for nonnormal data. Generalized linear mixed models (GLMM) will be used for multivariate analyses.^{120,121} The GLMM allow assessment of differences in Phase 1 versus Phase 3 for outcome variables that are binary, categorical, or continuous. Compared to logistic, multinomial, and multiple regression models, GLMM have the added advantage of simultaneously adjusting for the within-agency cluster correlation and the resulting increased variance.^{120,122} Continuous covariates will be tested for linearity and, if not linear, they will be transformed for linear fit using fractional polynomials or converted into categorical variables as appropriate.

The primary outcome variable (mortality), which is binary, will be modeled specifying a binomial distribution and logistic link function. Continuous outcome variables will use the GLMM with a Gaussian distribution and identity link function. The α -level is 0.05 for all statistical tests except the sequential analyses of the effect of implementation on mortality (interim, $\alpha = 0.01$; final, $\alpha = 0.04$). This gives an overall $\alpha = 0.05$ for the mortality analysis ($0.01 + 0.04 = 0.05$). All tests are two-tailed unless specified otherwise.

Analysis of Mortality. Because controlling for injury severity is essential when assessing the effect of an intervention on trauma survival,^{96,123–127} multiple complementary methodologies will be incorporated into the GLMM for this purpose. First, both the initial EMS and the initial trauma center Revised Trauma Scores (RTS) will be used in separate calculations. This is critical because guideline-based interventions may improve physiologic parameters that affect RTS (e.g., sBP). If this happens, then using only the trauma center RTS could be incorrectly interpreted as meaning that injury severity is lower in Phase 3 when, in fact, it was the EMS care that actually caused improved trauma center RTS values (and led to concomitant decreased mortality). Second, Injury Severity Score (ISS), a purely anatomic risk adjustment measure not influenced by treatment, will be used in the GLMM. In addition, a propensity score will be used to adjust for multiple severity-related parameters that are not included in the RTS or ISS (e.g., serum alcohol level).^{85,104,115,128–132} The propensity score will be included as a covariate in the GLMM. Given the data available in the ASTR (Data Supplement S2), multiple demographic (e.g., insurance status, race), medical (e.g., surgical interventions), and trauma center-associated variables (e.g., length of ICU stay) will be incorporated into the propensity score.^{104,130} A final propensity score will be generated by first incorporating all potentially relevant components and then reducing the list of components to optimally satisfy the balancing property.¹³² This will then be assessed for optimal fit in the

models by using both continuous and categorical versions.¹⁰⁴ A detailed statistical analysis plan is given in Data Supplements S3 and S4.

TRISS Methodology. The Trauma and Injury Severity Score (TRISS) methodology will be used to determine if the case severity has changed over time (Phase 1 vs. Phase 3). It is based on logistic regression coefficients derived from the Major Trauma Outcome Study (MTOS).^{123,124} TRISS has been validated in both adults and children and is used widely for risk adjustment in trauma.^{96,123–125,132–135} This will be used for stratification of survival across categories of survival probabilities and to assess for any differences in severity between study cohorts, comparing them to the reference group of the MTOS (details in Data Supplement S5, available as supporting information in the online version of this paper).

Analysis of Nonmortality (Secondary) Outcomes. Previous in-hospital reports have shown improvement in nonmortality outcomes from guideline therapy.^{86–89} The effect of EMS implementation on hospital/ICU length-of-stay, ventilator days, disposition (e.g., home, skilled nursing facility), and hospital cost (charges) will be compared in Phase 1 versus Phase 3 using the appropriate GLMM.

Analysis of Variations in Implementation. In a study of this size and diversity, variations in guideline utilization are inevitable.^{92,95,96,100–102,136} To test the effect of these variations, we will include interaction terms in the GLMM between variables related to guideline compliance. The fact that there will be “granularity” (e.g., variation in compliance of treating hypotension) will allow stratification of patients for subgroup analysis.

Analysis of ETCO₂ in Intubated Patients. Many animal and clinical studies have shown a profound negative effect of hyperventilation on cerebral blood flow and survival in TBI.^{8,10,14,17,50–53,55–76,85,104} Because of the large number of intubated cases, EPIC will allow specific analysis of the association between outcomes and 1) various ETCO₂ ranges and 2) the presence of ETCO₂ monitoring (~30% of agencies do not have monitors). Since the availability of ETCO₂ data will vary, several approaches will be considered, including ETCO₂ in the GLMMs as a predictor variable. The first approach will create categorical variables: none (no ETCO₂ data), normocapnea, single hypocapneic, multiple hypocapneic, single hypercapneic, and multiple hypercapneic episodes. Another approach will be to analyze the subgroups that have multiple or continuous measurements by regressing each individual’s ETCO₂ values versus time. This will allow us to determine if temporal trends in ETCO₂ are significant predictors of outcomes.

Analysis of Missing or Incomplete Data. Inevitably, some cases will have missing data for important covariates (although preliminary evaluation of the EMS data abstraction process reveals a very low missing data rate). Rather than removing these cases from analysis, multiple imputation will be used for missing covariates.^{93,115,135,137} Imputation of missing data is generally

preferable to alternatives, which are more likely to lead to biased and misleading results.^{93,115,137}

LIMITATIONS

Use of Diagnostic Criteria for Study Inclusion

No studies have evaluated which patient populations should receive guideline treatment. Thus, EPIC training emphasizes implementing the protocols in patients who have mechanisms consistent with significant TBI and who had loss of consciousness. The rationale for this broadly applied protocol was that we felt it was better to have some patients who received guideline treatment, but do not end up having major TBI, than to have patients who end up having major TBI, but did not receive guideline-based treatment. This approach is consistent with EPIC being a public health initiative. However, for the purpose of inclusion in the analysis, the patients must meet diagnostic and anatomic criteria for major TBI. This is important in a before–after study because it will prevent introduction of ascertainment bias by EMS personnel being more (or less) likely to “diagnose” TBI since the EMS impression will not determine inclusion in the analysis.

We believe that one of the strengths of EPIC is that it will allow, for the first time on such a large scale, the matching of data about EMS presentation and distal risk adjustment and outcome measures in true TBI. Hopefully this will help identify the patients who will most likely benefit from guideline-based treatment, based on their clinical presentations.

Nonrandomization

While no studies have evaluated the prehospital TBI guidelines, the supporting evidence is sufficient to have led to officially vetted guidelines.^{6,7,10,19,86,87,89} Thus, randomization (to guideline vs. nonguideline therapy) is probably not feasible since few, if any, systems would approve of randomization after full disclosure of the evidence. Because of this, we believe that the best available methodology to answer the critical questions proposed in this study is a large, prospective, before–after, observational trial.^{90,91,95–97,102,138,139}

Variations in Hospital Care

Interpretation of observational EMS trials can be complicated by changes in trauma center care. These may affect outcomes and can be inappropriately attributed to EMS interventions. To mitigate this potential we will conduct appropriate risk adjustment using factors such as diagnoses, ICD codes, probability of survival (TRISS),^{124,125,132–135} injury severity (e.g., ISS),^{124,125} and trauma center care.

Secular Trends in Severity

Coincidental changes in severity can affect outcomes. To evaluate this potential, we will perform concomitant analysis of patients transported by nonparticipating agencies to trauma centers. While we will not have detailed EMS data for these patients, we will have ASTR data. This will allow analysis for secular trends in demographics, injury diagnosis/severity, treatment, and outcomes, thus serving as a concurrent “control.”

Hawthorne Effect

During Phase 1 (before implementation), any potential differences between the retrospective and prospective groups will be assessed. This will yield useful information about the potential for an observer/Hawthorne effect that might occur simply because it is known that data are being collected.

CONCLUSIONS

The societal burden of traumatic brain injury is immense. While the potential for reducing morbidity and mortality by early treatment appears to be great, the effectiveness of the emergency medical services guidelines remains unproven. The Excellence in Prehospital Injury Care study will evaluate the effect of implementing the traumatic brain injury guidelines across a vast network of emergency medical services systems with an estimated enrollment of approximately 25,000 patients over the course of the study. Demonstrating the effect of guideline therapy would potentially lead to widespread implementation of the effective interventions. This could dramatically reduce morbidity and mortality from this major public health problem.

References

- Langlois JA, Rutland-Brown W, Wald MM. The epidemiology and impact of traumatic brain injury: a brief overview. *J Head Trauma Rehabil* 2006;21:375–8.
- Finkelstein E, Corso PS, Miller TR. *The Incidence and Economic Burden of Injuries in the United States*. Oxford, UK: Oxford University Press, 2006.
- Faul MD, Wald MM, Xu L, Coronado VG. *Traumatic Brain Injury in the United States: Emergency Department Visits, Hospitalizations and Deaths 2002–2006*. Available at: http://www.cdc.gov/traumaticbraininjury/pdf/blue_book.pdf. Accessed Apr 18, 2014.
- Thurman DJ, Alverson C, Dunn KA, Guerrero J, Sniezek JE. Traumatic brain injury in the United States: a public health perspective. *J Head Trauma Rehabil* 1999;14:602–15.
- Tepas JJ 3rd, DiScala C, Ramenofsky ML, Barlow B. Mortality and head injury: the pediatric perspective. *J Pediatr Surg* 1990;25:92–5.
- Adelson PD, Bratton SL, Carney NA, et al. Guidelines for the acute medical management of severe traumatic brain injury in infants, children, and adolescents. *Pediatr Crit Care Med* 2003;3(Suppl 1):S1–82.
- Brain Trauma Foundation, American Association of Neurological Surgeons, Congress of Neurological Surgeons, et al. Guidelines for the management of severe traumatic brain injury. *J Neurotrauma* 2007;24(Suppl 1):S1–106.
- Chesnut RM, Marshall LF, Klauber MR, et al. The role of secondary brain injury in determining outcome from severe head injury. *J Trauma* 1993;34:216–22.
- Jankowitz BT, Adelson PD. Pediatric traumatic brain injury: past, present and future. *Dev Neurosci* 2006;28:264–75.
- Badjatia N, Carney N, Crocco TJ, et al. Guidelines for prehospital management of traumatic brain injury, 2nd edition. *Prehosp Emerg Care* 2008;12(Suppl 1):S1–52.
- Bernard SA, Nguyen V, Cameron P, et al. Prehospital rapid sequence intubation improves functional outcome for patients with severe traumatic brain injury: a randomized controlled trial. *Ann Surg* 2010;252:959–65.
- Bulger EM, Copass MK, Sabath DR, Maier RV, Jurkovich GJ. The use of neuromuscular blocking agents to facilitate prehospital intubation does not impair outcome after traumatic brain injury. *J Trauma* 2005;58:718–23.
- Davis DP, Idris AH, Sise MJ, et al. Early ventilation and outcome in patients with moderate to severe traumatic brain injury. *Crit Care Med* 2006;34:1202–8.
- Davis DP, Dunford JV, Ochs M, Park K, Hoyt DB. The use of quantitative end-tidal capnometry to avoid inadvertent severe hyperventilation in patients with head injury after paramedic rapid sequence intubation. *J Trauma* 2004;56:808–14.
- Lewis F. Supply-dependent Oxygen Consumption: Reversing Cause and Effect. American Heart Association Resuscitation Science Symposium. New Orleans, LA, Nov 8, 2008.
- Davis DP, Fakhry SM, Wang HE, et al. Paramedic rapid sequence intubation for severe traumatic brain injury: perspectives from an expert panel. *Prehosp Emerg Care* 2007;11:1–8.
- Davis DP, Peay J, Serrano JA, et al. The impact of aeromedical response to patients with moderate to severe traumatic brain injury. *Ann Emerg Med* 2005;46:115–22.
- Eckstein M, Chan L, Schneir A, Palmer R. Effect of prehospital advanced life support on outcomes of major trauma patients. *J Trauma* 2000;48:643–8.
- Kochanek PM, Carney N, Adelson PD, et al. Guidelines for the acute medical management of severe traumatic brain injury in infants, children, and adolescents—second edition. *Pediatr Crit Care Med* 2012;13(Suppl 1):S1–82.
- Fearnside MR, Cook RJ, McDougall P, McNeil RJ. The Westmead Head Injury Project outcome in severe head injury. A comparative analysis of prehospital, clinical and CT variables. *Br J Neurosurg* 1993;7:267–79.
- Shutter LA, Narayan RK. Blood pressure management in traumatic brain injury [editorial]. *Ann Emerg Med* 2008;51(3 Suppl):S37–8.
- Gentleman D. Causes and effects of systemic complications among severely head injured patients transferred to a neurosurgical unit. *Int Surg* 1992;77:297–302.
- Pigula FA, Wald SL, Shackford SR, Vane DW. The effect of hypotension and hypoxia on children with severe head injuries. *J Pediatr Surg* 1993;28:310–4.
- Kokoska ER, Smith GS, Pittman T, Weber TR. Early hypotension worsens neurological outcome in pediatric patients with moderately severe head trauma. *J Pediatr Surg* 1998;33:333–8.

25. Miller JD, Becker DP. Secondary insults to the injured brain. *J R Coll Surg Edinb* 1982;27:292–8.
26. Barton CW, Hemphill JC, Morabito D, Manley G. A novel method of evaluating the impact of secondary brain insults on functional outcomes in traumatic brain-injured patients. *Acad Emerg Med* 2005;12:1–6.
27. Manley G, Knudson MM, Morabito D, Damron S, Erickson V, Pitts L. Hypotension, hypoxia, and head injury: frequency, duration, and consequences. *Arch Surg* 2001;136:1118–23.
28. Mayer TA, Walker ML. Pediatric head injury: the critical role of the emergency physician. *Ann Emerg Med* 1985;14:1178–84.
29. Ong L, Selladurai BM, Dhillon MK, Atan M, Lye MS. The prognostic value of the Glasgow Coma Scale, hypoxia and computerised tomography in outcome prediction of pediatric head injury. *Pediatr Neurosurg* 1996;24:285–91.
30. Price DJ, Murray A. The influence of hypoxia and hypotension on recovery from head injury. *Injury* 1972;3:218–24.
31. Stocchetti N, Furlan A, Volta F. Hypoxemia and arterial hypotension at the accident scene in head injury. *J Trauma* 1996;40:764–7.
32. Luerksen TG, Klauber MR, Marshall LF. Outcome from head injury related to patient's age. A longitudinal prospective study of adult and pediatric head injury. *J Neurosurg* 1988;68:409–16.
33. Carrel M, Moeschler O, Ravussin P, Favre JB, Boulard G. Prehospital air ambulance and systemic secondary cerebral damage in severe craniocerebral injuries. *Ann Fr Anesth Reanim* 1994;13:326–35.
34. Jeffreys RV, Jones JJ. Avoidable factors contributing to the death of head injury patients in general hospitals in Mersey Region. *Lancet* 1981;2:459–61.
35. Kohi YM, Mendelow AD, Teasdale GM, Allardice GM. Extracranial insults and outcome in patients with acute head injury—relationship to the Glasgow Coma Scale. *Injury* 1984;16:25–9.
36. Rose J, Valtonen S, Jennett B. Avoidable factors contributing to death after head injury. *Br Med J* 1977;2:615–8.
37. Seelig JM, Klauber MR, Toole BM, Marshall LF, Bowers SA. Increased ICP and systemic hypotension during the first 72 hours following severe head injury. In: Miller JD, Teasdale GM, Rowan JO (editors). *Intracranial Pressure VI*. Berlin, Germany: Springer-Verlag, 1986, pp 675–9.
38. Chesnut RM, Ghajar J, Maas AIR, et al. Guidelines for the management of severe traumatic brain injury. Part 2: early indicators of prognosis in severe traumatic brain injury. *J Neurotrauma* 2000;17:555–627.
39. Hill DA, Abraham KJ, West RH. Factors affecting outcome in the resuscitation of severely injured patients. *Aust N Z J Surg* 1993;63:604–9.
40. Davis DP, Dunford JV, Poste JC, et al. The impact of hypoxia and hyperventilation on outcome after paramedic rapid sequence intubation of severely head-injured patients. *J Trauma* 2004;57:1–8.
41. Dunford JV, Davis DP, Ochs M, Doney M, Hoyt DB. Incidence of transient hypoxia and pulse rate reactivity during paramedic rapid sequence intubation. *Ann Emerg Med* 2003;42:721–8.
42. Jones PA, Andrews PJ, Midgley S, et al. Measuring the burden of secondary insults in head-injured patients during intensive care. *J Neurosurg Anesthesiol* 1994;6:4–14.
43. Marmarou A, Anderson RL, Ward JD, et al. Impact of ICP instability and hypotension on outcome in patients with severe head trauma. *J Neurosurg* 1991;75(Suppl):S59–S66.
44. Silverston P. Pulse oximetry at the roadside: a study of pulse oximetry in immediate care. *Br Med J* 1989;298:711–3.
45. Darby JM, Yonas H, Marion DW, Latchaw RE. Local “inverse steal” induced by hyperventilation in head injury. *Neurosurgery* 1988;23:84–8.
46. Gordon E, Rossanda M. The importance of the cerebrospinal fluid acid-base status in the treatment of unconscious patients with brain lesions. *Acta Anaesthesiol Scand* 1968;12:51–73.
47. Obrist WD, Clifton GL, Robertson CS. Cerebral metabolic changes induced by hyperventilation in acute head injury. In: Meyer JS (editor). *Cerebral Vascular Disease 6*. Philadelphia, PA: Elsevier Science, 1987, pp 241–55.
48. Bruce DA, Raphaely RC, Goldberg AI, et al. Pathophysiology, treatment and outcome following severe head-injury in children. *Childs Brain* 1979;5:174–91.
49. Raphaely RC, Swedlow DB, Downes JJ, Bruce DA. Management of severe pediatric head trauma. *Pediatr Clin North Am* 1980;27:715–27.
50. Muizelaar JP, Marmarou A, Ward JD, et al. Adverse effects of prolonged hyperventilation in patients with severe head injury: a randomized clinical trial. *J Neurosurg* 1991;75:731–9.
51. Davis DP. Early ventilation in traumatic brain injury. *Resuscitation* 2008;76:333–40.
52. Davis DP, Heister R, Poste JC, Hoyt DB, Ochs M, Dunford JV. Ventilation patterns in patients with severe traumatic brain injury following paramedic rapid sequence intubation. *Neurocrit Care* 2005;2:165–71.
53. Manley GT, Hemphill JC, Morabito D, et al. Cerebral oxygenation during hemorrhagic shock: perils of hyperventilation and the therapeutic potential of hypoventilation. *J Trauma* 2000;48:1025–32.
54. Wang HE, Davis DP, O'Connor RE, Domeier RM. Drug-assisted intubation in the prehospital setting (resource document to NAEMSP position statement). *Prehosp Emerg Care* 2006;10:261–71.
55. Pepe PE, Lurie KG, Wigginton JG, Raedler C, Idris AH. Detrimental hemodynamic effects of assisted ventilation in hemorrhagic states. *Crit Care Med* 2004;32(9 Suppl):S414–20.
56. Pepe PE, Roppolo LP, Fowler RL. The detrimental effects of ventilation during low-blood-flow states. *Curr Opin Crit Care* 2005;11:212–8.
57. Poste JC, Davis DP, Ochs M, et al. Air medical transport of severely head-injured patients undergoing

- paramedic rapid sequence intubation. *Air Med J* 2004;23:36–40.
58. Prause G, Hetz H, Lauda P, Pojer H, Smolle-Juettner F, Smolle J. A comparison of the end-tidal CO₂ documented by capnometry and the arterial pCO₂ in emergency patients. *Resuscitation* 1997;35:145–8.
 59. Robertson CS, Valadka AB, Hannay HJ, et al. Prevention of secondary ischemic insults after severe head injury. *Crit Care Med* 1999;27:2086–95.
 60. Bao Y, Jiang J, Zhu C, Lu Y, Cai R, Ma C. Effect of hyperventilation on brain tissue oxygen pressure, carbon dioxide pressure, pH value and intracranial pressure during intracranial hypertension in pigs. *Chin J Traumatol* 2000;3:210–3.
 61. Forbes ML, Clark RS, Dixon CE, et al. Augmented neuronal death in CA3 hippocampus following hyperventilation early after controlled cortical impact. *J Neurosurg* 1998;88:549–56.
 62. Skippen P, Seear M, Poskitt K, et al. Effect of hyperventilation on regional cerebral blood flow in head-injured children. *Crit Care Med* 1997;25:1402–9.
 63. Coles JP, Minhas PS, Fryer TD, et al. Effect of hyperventilation on cerebral blood flow in traumatic head injury: clinical relevance and monitoring correlates. *Crit Care Med* 2002;30:1950–9.
 64. Ausina A, Báguena M, Nadal M, et al. Cerebral hemodynamic changes during sustained hypocapnia in severe head injury: can hyperventilation cause cerebral ischemia? *Acta Neurochir Suppl* 1998;71:1–4.
 65. Marion DW, Puccio A, Wisniewski SR, et al. Effect of hyperventilation on extracellular concentrations of glutamate, lactate, pyruvate, and local cerebral blood flow in patients with severe traumatic brain injury. *Crit Care Med* 2002;30:2619–25.
 66. Adelson PD, Clyde B, Kochanek PM, Wisniewski SR, Marion DW, Yonas H. Cerebrovascular response in infants and young children following severe traumatic brain injury: a preliminary report. *Pediatr Neurosurg* 1997;26:200–7.
 67. Kiening KL, Hartl R, Unterberg AW, Schneider GH, Bardt T, Lanksch WR. Brain tissue pO₂-monitoring in comatose patients: implications for therapy. *Neurol Res* 1997;19:233–40.
 68. Sharples PM, Matthews DS, Eyre JA. Cerebral blood flow and metabolism in children with severe head injuries. Part 2: cerebrovascular resistance and its determinants. *J Neurol Neurosurg Psychiatry* 1995;58:153–9.
 69. Muizelaar JP, Marmarou A, DeSalles AA, et al. Cerebral blood flow and metabolism in severely head-injured children. Part 1: relationship with GCS score, outcome, ICP, and PVI. *J Neurosurg* 1989;71:63–71.
 70. Bouma GJ, Muizelaar JP, Choi SC, Newlon PG, Young HF. Cerebral-circulation and metabolism after severe traumatic brain injury—the elusive role of ischemia. *J Neurosurg* 1991;75:685–93.
 71. Bouma GJ, Muizelaar JP, Stringer WA, Choi SC, Fatouros P, Young HF. Ultra-early evaluation of regional cerebral blood-flow in severely head-injured patients using xenon-enhanced computerized-tomography. *J Neurosurg* 1992;77:360–8.
 72. Fieschi C, Battistini N, Beduschi A, Boselli L, Rossanda M. Regional cerebral blood flow and intraventricular pressure in acute head injuries. *J Neurol Neurosurg Psychiatry* 1974;37:1378–88.
 73. Jaggi JL, Obrist WD, Gennarelli TA, Langfitt TW. Relationship of early cerebral blood flow and metabolism to outcome in acute head injury. *J Neurosurg* 1990;72:176–82.
 74. Marion DW, Darby J, Yonas H. Acute regional cerebral blood flow changes caused by severe head injuries. *J Neurosurg* 1991;74:407–14.
 75. Fortune JB, Feustel PJ, Graca L, Hasselbarth J, Kuehler DH. Effect of hyperventilation, mannitol, and ventriculostomy drainage on cerebral blood flow after head injury. *J Trauma* 1995;39:1091–7.
 76. Sioutos PJ, Orozco JA, Carter LP, Weinand ME, Hamilton AJ, Williams FC. Continuous regional cerebral cortical blood flow monitoring in head-injured patients. *Neurosurgery* 1995;36:943–9.
 77. Davis DP, Douglas DJ, Koenig W, Carrison D, Buono C, Dunford JV. Hyperventilation following aero-medical rapid sequence intubation may be a deliberate response to hypoxemia. *Resuscitation* 2007;73:354–61.
 78. Thomas SH, Orf J, Wedel SK, Conn AK. Hyperventilation in traumatic brain injury patients: inconsistency between consensus guidelines and clinical practice. *J Trauma* 2002;52:47–52.
 79. Aufderheide TP, Lurie KG. Death by hyperventilation: a common and life-threatening problem during cardiopulmonary resuscitation. *Crit Care Med* 2004;32(9 Suppl):S345–51.
 80. Braman SS, Dunn SM, Amico CA, Millman RP. Complications of intrahospital transport in critically ill patients. *Ann Intern Med* 1987;107:469–73.
 81. Gervais HW, Eberle B, Konietzke D, Hennes HJ, Dick W. Comparison of blood gases of ventilated patients during transport. *Crit Care Med* 1987;15:761–3.
 82. Helm M, Hauke J, Lampl L. A prospective study of the quality of pre-hospital emergency ventilation in patients with severe head injury. *Br J Anaesth* 2002;88:345–9.
 83. Hurst JM, Davis K Jr, Branson RD, Johannigman JA. Comparison of blood gases during transport using two methods of ventilatory support. *J Trauma* 1989;29:1637–40.
 84. Tobias JD, Lynch A, Garrett J. Alterations of end-tidal carbon dioxide during the intrahospital transport of children. *Pediatr Emerg Care* 1996;12:249–51.
 85. Denninghoff KR, Griffin MJ, Bartolucci AA, Lobello SG, Fine PR. Emergent endotracheal intubation and mortality in traumatic brain injury. *West J Emerg Med* 2008;9:184–9.
 86. Faul M, Wald MM, Rutland-Brown W, Sullivent EE, Sattin RW. Using a cost-benefit analysis to estimate outcomes of a clinical treatment guideline: testing the Brain Trauma Foundation guidelines for the treatment of severe traumatic brain injury. *J Trauma* 2007;63:1271–8.

87. Palmer S, Bader MK, Qureshi A, et al. The impact on outcomes in a community hospital setting of using the AANS traumatic brain injury guidelines. *American Association of Neurological Surgeons. J Trauma* 2001;50:657–64.
88. Kreutzer JS, Kolakowsky-Hayner SA, Ripley D, et al. Charges and lengths of stay for acute and inpatient rehabilitation treatment of traumatic brain injury 1990–1996. *Brain Inj* 2001;15:763–74.
89. Fakhry SM, Trask AL, Waller MA, Watts DD. Management of brain-injured patients by an evidence-based medicine protocol improves outcomes and decreases hospital charges. *J Trauma* 2004;56:492–9.
90. Keim SM, Spaite DW, Maio RF, et al. Establishing the scope and methodological approach to out-of-hospital outcomes and effectiveness research. *Acad Emerg Med* 2004;11:1067–73.
91. Spaite DW, Maio R, Garrison HG, et al. Emergency Medical Services Outcomes Project (EMSOP) II: developing the foundation and conceptual models for out-of-hospital outcomes research. *Ann Emerg Med* 2001;37:657–63.
92. Bobrow BJ, Clark LL, Ewy GA, et al. Minimally interrupted cardiac resuscitation by emergency medical services for out-of-hospital cardiac arrest. *JAMA* 2008;299:1158–65.
93. Bobrow BJ, Spaite DW, Berg RA, et al. Chest compression-only CPR by lay rescuers and survival from out-of-hospital cardiac arrest. *JAMA* 2010;304:1447–54.
94. Spaite DW, Beskind DL, Garrison HG. Evaluating the effectiveness of EMS systems: utilizing outcomes research methods to identify the impact of prehospital care. In: Cone DC, O’Conner RE, Fowler R (editors). *Emergency Medical Services: Clinical Practice and Systems Oversight*. Lenexa, KS: National Association of EMS Physicians, 2009.
95. Stiell IG, Wells GA, Spaite DW, et al. The Ontario Prehospital Advanced Life Support (OPALS) study part II: rationale and methodology for trauma and respiratory distress patients. OPALS Study Group. *Ann Emerg Med* 1999;34:256–62.
96. Stiell IG, Nesbitt LP, Pickett W, et al. The OPALS Major Trauma Study: impact of advanced life-support on survival and morbidity. *CMAJ* 2008;178:1141–52.
97. Stiell IG, Spaite DW, Field B, et al. Advanced life support for out-of-hospital respiratory distress. *N Engl J Med* 2007;356:2156–64.
98. Stiell IG, Wells GA, Field BJ, et al. Improved out-of-hospital cardiac arrest survival through the inexpensive optimization of an existing defibrillation program: OPALS study phase II. *Ontario Prehospital Advanced Life Support. JAMA* 1999;281:1175–81.
99. Stiell IG, Wells GA, Field B, et al. Advanced cardiac life support in out-of-hospital cardiac arrest. *N Engl J Med* 2004;351:647–56.
100. Bobrow BJ, Ewy GA, Clark L, et al. Passive oxygen insufflation is superior to bag-valve-mask ventilation for witnessed ventricular fibrillation out-of-hospital cardiac arrest. *Ann Emerg Med* 2009;54:656–62.
101. Bobrow BJ, Vadeboncoeur TF, Clark L, Chikani V. Establishing Arizona’s statewide cardiac arrest reporting and educational network. *Prehosp Emerg Care* 2008;12:381–7.
102. Spaite DW, Bobrow BJ, Vadeboncoeur TF, et al. The impact of prehospital transport interval on survival in out-of-hospital cardiac arrest: implications for regionalization of post-resuscitation care. *Resuscitation* 2008;79:61–6.
103. Maas AI, Stocchetti N, Bullock R. Moderate and severe traumatic brain injury in adults. *Lancet Neurol* 2008;7:728–41.
104. Wang HE, Peitzman AB, Cassidy LD, Adelson PD, Yealy DM. Out-of-hospital endotracheal intubation and outcome after traumatic brain injury. *Ann Emerg Med* 2004;44:439–50.
105. Barell V, Aharonson-Daniel L, Fingerhut LA, et al. An introduction to the Barell body region by nature of injury diagnosis matrix. *Inj Prev* 2002;8:91–6.
106. Clark DE, Ahmad S. Estimating injury severity using the Barell matrix. *Inj Prev* 2006;12:111–6.
107. Fingerhut LA, Aharonson-Daniel L, Mackenzie EJ, et al. The Barell matrix. *Inj Prev* 2002;8:259.
108. Iezzoni LI. *Risk Adjustment for Measuring Health Care Outcomes*, 3rd ed. Chicago, IL: Health Administration Press, 2003.
109. Garrison HG, Maio RF, Spaite DW, et al. Emergency Medical Services Outcomes Project III (EMSOP III): the role of risk adjustment in out-of-hospital outcomes research. *Ann Emerg Med* 2002;40:79–88.
110. Hsia RY, Wang E, Torres H, Saynina O, Wise PH. Disparities in trauma center access despite increasing utilization: data from California, 1999 to 2006. *J Trauma* 2010;68:217–24.
111. Newgard CD, Zive D, Holmes JF, et al. A multisite assessment of the American College of Surgeons Committee on Trauma field triage decision scheme for identifying seriously injured children and adults. *J Am Coll Surg* 2011;213:709–21.
112. Nakamura Y, Daya M, Bulger EM, et al. Evaluating age in the field triage of injured persons. *Ann Emerg Med* 2012;60:335–45.
113. McPherson K. Sequential stopping rules in clinical trials. *Stat Med* 1990;9:595–600.
114. O’Brien PC, Fleming TR. A multiple testing procedure for clinical trials. *Biometrics* 1979;35:549–56.
115. Pocock SJ. Interim analysis and stopping rules for clinical trials. In: Bithell JF, Coppi R (editors). *Perspectives in Medical Statistics*. Philadelphia, PA: Academic Press, Elsevier, 1981.
116. Armitage P. *Sequential Medical Trials*. 2nd ed. Oxford, UK: Blackwell Scientific Publications, 1975.
117. Peto R, Pike MC, Armitage P, et al. Design and analysis of randomized clinical trials requiring prolonged observation of each patient. I. Introduction and design. *Br J Cancer* 1976;34:585–612.
118. Erdfelder E, Faul F, Buchner A. GPOWER: a general power analysis program. *Behav Res Methods Instr Comput* 1996;28:1–11.
119. Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: a flexible statistical power analysis

- program for the social, behavioral, and biomedical sciences. *Behav Res Methods* 2007;39:175–91.
120. McCulloch CE, Searle SR. *Generalized, Linear and Mixed Models*. New York, NY: Wiley, 2001.
 121. McCullagh P, Nelder JA. *Generalized Linear Models*. 2nd ed. London, UK: Chapman & Hall, 1989.
 122. Karim MR, Zeger SL. Generalized linear models with random effects; salamander mating revisited. *Biometrics* 1992;48:631–44.
 123. Champion HR, Sacco WJ, Copes WS. Injury severity scoring again. *J Trauma* 1995;38:94–5.
 124. Champion HR, Copes WS, Sacco WJ, et al. The Major Trauma Outcome Study: establishing national norms for trauma care. *J Trauma* 1990;30:1356–65.
 125. Champion HR. Trauma scoring. *Scand J Surg* 2002;91:12–22.
 126. Senkowski CK, McKenney MG. Trauma scoring systems: a review. *J Am Coll Surg* 1999;189:491–503.
 127. MacKenzie EJ, Rivara FP, Jurkovich GJ, et al. A national evaluation of the effect of trauma-center care on mortality. *N Engl J Med* 2006;354:366–78.
 128. Newgard CD, Hedges JR, Arthur M, Mullins RJ. Advanced statistics: the propensity score—a method for estimating treatment effect in observational research. *Acad Emerg Med* 2004;11:953–61.
 129. D’Agostino RB Jr. Propensity score methods for bias reduction in the comparison of a treatment to a non-randomized control group. *Stat Med* 1998;17:2265–81.
 130. Cudnik MT, Newgard CD, Daya M, Jui J. The impact of rapid sequence intubation on trauma patient mortality in attempted prehospital intubation. *J Emerg Med* 2010;38:175–81.
 131. Rubin DB. Estimating causal effects from large data sets using propensity scores. *Ann Intern Med* 1997;127:757–63.
 132. Marcin JP, Pollack MM. Triage scoring systems, severity of illness measures, and mortality prediction models in pediatric trauma. *Crit Care Med* 2002;30:S457–67.
 133. Boyd CR, Tolson MA, Copes WS. Evaluating trauma care: the TRISS method. Trauma Score and the Injury Severity Score. *J Trauma* 1987;27:370–8.
 134. Ott R, Kramer R, Martus P, Bussenius-Kammerer M, Carbon R, Rupperecht H. Prognostic value of trauma scores in pediatric patients with multiple injuries. *J Trauma* 2000;49:729–36.
 135. Kaufmann CR, Maier RV, Kaufmann EJ, Rivara FP, Carrico CJ. Validity of applying adult TRISS analysis to injured children. *J Trauma* 1991;31:691–7.
 136. Spaite DW, Criss EA, Valenzuela TD, Guisto J. Emergency medical service systems research: problems of the past, challenges of the future. *Ann Emerg Med* 1995;26:146–52.
 137. Bobrow BJ, Vadeboncoeur TF, Stolz U, et al. The influence of scenario-based training and real-time audiovisual feedback on out-of-hospital cardiopulmonary resuscitation quality and survival from out-of-hospital cardiac arrest. *Ann Emerg Med* 2013;62:47–56.
 138. Engdahl J, Abrahamsson P, Bang A, et al. Is hospital care of major importance for outcome after out-of-hospital cardiac arrest? Experience acquired from patients with out-of-hospital cardiac arrest resuscitated by the same Emergency Medical Service and admitted to one of two hospitals over a 16-year period in the municipality of Goteborg. *Resuscitation* 2000;43:201–11.
 139. McLean SA, Maio RF, Spaite DW, Garrison HG. Emergency medical services outcomes research: evaluating the effectiveness of prehospital care. *Prehosp Emerg Care* 2002;6:S52–6.
 140. Gausche M, Lewis RJ, Stratton SJ, et al. Effect of out-of-hospital pediatric endotracheal intubation on survival and neurological outcome: a controlled clinical trial. *JAMA* 2000;283:783–90.
 141. Warner KJ, Cuschieri J, Copass MK, Jurkovich GJ, Bulger EM. The impact of prehospital ventilation on outcome after severe traumatic brain injury. *J Trauma* 2007;62:1330–6.
 142. Warner KJ, Bulger EM. Does pre-hospital ventilation effect outcome after significant brain injury. *Trauma* 2007;9:283–89.
 143. Brower RG, Matthay MA, Morris A, et al. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med* 2000;342:1301–8.
 144. Bloomfield GL, Ridings PC, Blocher CR, Marmarou A, Sugerman HJ. A proposed relationship between increased intraabdominal, intrathoracic, and intracranial pressure. *Crit Care Med* 1997;25:496–503.
 145. Citerio G, Vascotto E, Villa F, Celotti S, Pesenti A. Induced abdominal compartment syndrome increases intracranial pressure in neurotrauma patients: a prospective study. *Crit Care Med* 2001;29:1466–71.
 146. Slutsky AS, Ranieri VM. Mechanical ventilation: lessons from the ARDSNet trial. *Respir Res* 2000;1:73–7.
 147. Uhlig S. Ventilation-induced lung injury and mechanotransduction: stretching it too far? *Am J Physiol Lung Cell Mol Physiol* 2002;282:L892–6.
 148. American Heart Association. Pediatric Advanced Life Support (PALS). Available at: http://www.heart.org/HEARTORG/CPRandECC/HealthcareProviders/Pediatrics/Pediatric-Advanced-Life-Support-PALS_UCM_303705_Article.jsp. Accessed Apr 13, 2014.
 149. Helmy A, Vizcaychipi M, Gupta AK. Traumatic brain injury: intensive care management. *Br J Anaesth* 2007;99:32–42.
 150. Spaite DW, Valenzuela TD, Meislin HW, Criss EA, Hinsberg P. Prospective validation of a new model for evaluating emergency medical services systems by in-field observation of specific time intervals in prehospital care. *Ann Emerg Med* 1993;22:638–45.

Supporting Information

The following supporting information is available in the online version of this paper:

Data Supplement S1. EPIC EMS dataset.

Data Supplement S2. ASTR—required data elements for trauma center reporting.

Data Supplement S3. EPIC—adults (age \geq 18 years): analysis and statistical approach.

Data Supplement S4. EPIC4Kids (age $<$ 18 years): analysis and statistical approach.

Data Supplement S5. Use of TRISS methodology in the EPIC analysis.

Abstracts en español!

Beginning with the September issue, *Academic Emergency Medicine* will be publishing the abstracts of the various articles in Spanish. They will be presented alongside the English abstracts in the online versions of each paper (pdf, html, and mobile apps). The Spanish abstracts will also be included in the print edition of the journal for any papers that originate in Spanish-speaking countries, or are likely to be of particular interest to emergency physicians in Spanish-speaking countries.

This project would not be possible without technical assistance and generous funding from our publisher, John Wiley and Sons, Inc., and the language assistance of *Emergencias*, the journal of the Sociedad Española de Medicina de Urgencias y Emergencias (SEMES).